

Clinical Frailty Scale and Body Mass Index as an Independent Predictor of 2-year Mortality at Hospitalized Patients

Yelda Öztürk, Arzu Okyar Baş, Merve Güner, Serdar Ceylan, Meltem Koca, Mert Esmе, Cafer Balcı, Burcu Balam Doğu, Mustafa Cankurtaran, Meltem Halil

Hacettepe University Faculty of Medicine, Department of Internal Medicine, Division of Geriatrics, Ankara, Turkey

Abstract

Objective: This study aimed to investigate the effect of the clinical frailty scale (CFS) and body mass index (BMI) on the 2-year mortality prediction in hospitalized internal medicine patients.

Materials and Methods: A prospective observational study was conducted between January 2019 and February 2020. Subjects (18 years and older) admitted to the internal medicine wards and expected to stay for at least 72 h were included. Participants were evaluated within 48 h of admission. The Charlson comorbidity index (CCI) was calculated. Anthropometric measurements and handgrip strength were obtained within 48 h. CFS was used for frailty assessment. Cox regression analysis was performed for mortality analysis.

Results: One hundred eighteen patients were included. Fifty-eight of the (49.2%) patients were 65 years and over. In multivariate analysis, BMI and CFS were independently associated with 2-year mortality, regardless of age, sex, and CCI. Hazard ratios for BMI and CFS were 0.898 [95% confidence interval (CI), 0.840-0.961; $p=0.002$] and 1.313 (95% CI, 1.002-1.719; $p=0.048$), respectively.

Conclusion: Higher CFS scores and lower BMI scores are independently associated with 2-year mortality in hospitalized internal medicine patients.

Keywords: Body mass index, clinical frailty scale, frailty, hospitalization, mortality

Introduction

Frailty is a state of increased vulnerability to stressors and is also associated with multiple physiological systems that are interrelated with each other (1). It is a global public healthcare issue as the world is aging. Subjects living with frailty are at a growing risk of adverse outcomes, including hospitalization and mortality, causing higher healthcare costs (2). It is a known fact that subjects with frailty can dynamically transition between states (3). Therefore, it is crucial to detect and manage subjects who are living with frailty.

The prevalence of frailty in geriatric inpatients ranges from 48.8% to 80%, depending on the evaluation tool used (4). There are several frailty instruments such as the FRAIL scale, Edmonton frailty scale, and clinical frailty scale (CFS) (3,5-7). CFS is an

easy and quick scale. It was developed to determine frailty in older adults and includes items such as comorbidity, cognitive impairment, and function (8). It assesses frailty using visual and written charts with nine graded pictures, ranging between 1 (very fit) and 9 (terminally ill). A score of ≥ 5 represents patients who are frail. CFS has been shown to be widely used in multiple settings. Several studies have been conducted, especially in hospital settings, and assessed its associations with adverse outcomes (9).

Body mass index (BMI) is also known to be a factor related to mortality. It is an index of malnutrition (10). Malnutrition (both undernutrition and obesity) plays a key role in the pathogenesis of frailty and sarcopenia (11). A recently published meta-analysis revealed a high overlapping prevalence of malnutrition

Address for Correspondence: Yelda Öztürk, Eskişehir City Hospital, Clinic of Geriatrics, Eskişehir, Turkey

Phone: +90 533 615 12 86 **E-mail:** yeldaoszturk67@gmail.com **ORCID:** orcid.org/0000-0003-3847-0967

Received: 27.03.2023 **Accepted:** 16.08.2023

Cite this article as: Öztürk Y, Okyar Baş A, Güner M, Ceylan S, Koca M, Esmе M, Balcı C, Doğu BB, Cankurtaran M, Halil M. Clinical Frailty Scale and Body Mass Index as an Independent Predictor of 2-year Mortality at Hospitalized Patients. Eur J Geriatr Gerontol 2024;6(1):26-30



Copyright© 2024 The Author. Published by Galenos Publishing House on behalf of Turkish Academic Geriatrics Society.

This is an open access article under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND) International License.



and frailty in hospitalized older patients (12). On the other hand, obesity has a close relationship with type 2 diabetes mellitus and coronary artery disease (13). However, there are conflicting results regarding its effect on mortality, changing from condition to condition (14,15).

Obesity paradox, a term used to describe that overweight and obese patients with a particular disease may have better outcomes than normal weight patients, is another concern (16,17). For example, it has been claimed that the strength of the association between obesity and mortality weakens with increasing age (18). There is wide heterogeneity between studies regarding the relationship between obesity and mortality, especially in older patients (19).

In light of these data, we determined the effect of CFS and BMI on 2-year mortality in hospitalized internal medicine patients.

Materials and Methods

This prospective observational study was conducted between January 2019 and February 2020. Subjects (18 years and older), admitted to the internal medicine wards of a university hospital and expected to stay for at least 72 h, were included. Written informed consents were obtained from all participants. Participants were evaluated within 48 h of admission, and they were followed up for at least 2 years or until death. The baseline characteristics of patients, including comorbidities, were recorded. The Charlson comorbidity index (CCI) was calculated (20).

Anthropometric measurements including height, weight, calf circumference (CC), and mid-upper arm circumference (MAC) were taken. Height and weight were measured while standing and recorded in meters and kilograms, respectively (TEM-BEKO 035x040, İstanbul, Turkey). BMI was calculated by dividing body weight in kilograms by height in meters squared (kg/m^2). CC was measured in the sitting position with 90° of knee flexion at the largest level of the leg. MAC was measured when the elbow was at 90° flexion. CC and MAC were recorded in centimeters. Handgrip strength (HGS) was measured by Takei digital grip strength dynamometer (Takei Scientific Instruments Co, Niigata, Japan) at a 90° flexion of the elbow with a neutrally rotated forearm and reported in kilograms. The thresholds of 16 kg for females and 27 kg for males were used, as recommended by EWGSOP-2 (21). The highest value of the three measurements was considered (22). Mini nutrition assessment-short form (MNA-SF), and nutritional risk score-2002 (NRS-2002) were performed to screen for malnutrition. Patients were grouped as malnourished (score ≤ 7), at-risk (score 8–11), and normal (score 12–14) according to the MNA-SF score (23). When the NRS-2002 score was ≥ 3 , it meant nutritionally at risk (24). Frailty was assessed using the 9-point CFS. The score ranged from 1 to 9 (25). The study was approved by the Hacettepe University Non-

Interventional Clinical Research Ethics Committee (decision number: 2019/07-28, date: 07.03.2019).

Statistics

The IBM SPSS Statistics program version 23.0 was used for statistical analysis. The normality of variables was examined using visual (histograms and probability plots) and analytical methods. Categorical variables are presented as numbers and frequencies. Normally distributed variables are presented as mean \pm standard deviation, non-normally distributed variables are presented as median (IQR, 25p–75p). Patients were divided into two groups as younger (<65 years) and older (≥ 65 years) to present baseline characteristics of patients. The χ^2 or Fisher's exact test was used to compare differences between the categorical variables as appropriate. Mann-Whitney U and Student's t-tests were used to compare non-normally and normally distributed variables, respectively. Cox regression analysis was performed to define the factors associated with 2-year mortality. Four models were constructed. Model 1 included age and sex; model 2 included age, sex, and CCI; model 3 included age, sex, CCI, and BMI; model 4 included age, sex, CCI, BMI, and CFS. The findings are shown as hazard ratios (HRs) and the corresponding 95% confidence interval (CI). The proportional hazard assumption and model fit were assessed using residual (Schoenfeld and Martingale) analysis. All analyzes were considered statistically significant when the p value was <0.05 .

Results

A total of 118 patients were included in the analysis. Subjects were divided into two groups as younger ($n=60$) and older ($n=58$). The baseline characteristics of the patients are presented in Table 1. Causes of hospitalization, length of stay, CCI, MAC, CC, and MNA-SF categories were not different between the groups. Median BMI values of the younger and older groups were 25.8 (22.7–29.9), and 29.1 (25.5–32.1), respectively ($p=0.008$). The rate of patients with risk of malnutrition according to NRS-2002 was higher in the older group ($p=0.001$). The median CFS score was higher in the older group than the younger ($p>0.001$). The rates of patients with low muscle strength were 38.6% ($n=22$) in the younger group and 74.5% ($n=41$) in the older group ($p<0.001$).

During the 2-year follow-up, 28.8% of patients died (21.7% of younger group, 36.2% of older group). Age ($p=0.015$), CCI ($p=0.021$), BMI ($p=0.032$) and CFS ($p=0.001$) were significantly associated with 2-year mortality in the univariate model (Table 2). Four different models were created and are presented in Table 3. In model 4, which included age, sex, CCI, BMI, and CFS, BMI and CFS were independently associated with 2-year mortality. The HRs for BMI and CFS were 0.898 (95% CI, 0.840–0.961; $p=0.002$) and 1.313 (95% CI, 1.002–1.719; $p=0.048$), respectively.

Table 1. Baseline characteristics of younger and older patients

	<65 years (n=60)	≥65 years (n=58)	p
Sex, female	39 (65.0)	27 (46.6)	0.044
Illiterate	6 (10.0)	16 (27.6)	<0.001
Causes of hospitalization			
Malignancy	5 (8.3)	7 (12.1)	0.502
Infectious disease	5 (8.3)	7 (12.1)	0.502
Rheumatic disease	3 (5.0)	5 (8.6)	0.434
Gastrointestinal disease	7 (11.7)	10 (17.2)	0.389
Hematologic disease	16 (26.7)	5 (8.6)	0.010
Endocrine disease	8 (13.3)	2 (3.4)	0.054
Pulmonary disease	6 (10.0)	3 (5.2)	0.323
Nephrology disorders	5 (8.3)	11 (19.0)	0.092
Cardiovascular disease	5 (8.3)	8 (13.8)	0.344
Length of hospital stay, days	12.8 (8.8-19.7)	10.8 (7.8-17.0)	0.391
Charlson comorbidity index	2 (1-3)	2 (1-3)	0.191
Body mass index, kg/m ²	25.8 (22.7-29.9)	29.1 (25.5-32.1)	0.008
Mid-upper arm circumference, cm	28 (24-30)	27 (25-30)	0.606
Calf circumference, cm	34 (31-37)	35 (31-37)	0.793
NRS-2002, at risk (≥3)	13 (21.7)	30 (51.7)	0.001
MNA-SF score ≤11	28 (43.8)	36 (56.3)	0.093
Clinical frailty scale	3 (2-4)	4 (4-5)	<0.001
Low handgrip strength, kg	22 (38.6)	41 (74.5)	<0.001

Numbers are means ± SD, medians (25th-75th percentiles), or frequencies (%), as appropriate
MNA-SF: Mini nutrition assessment-short form, NRS-2002: Nutritional risk score-2002, SD: Standard deviation

Table 2. Univariable analysis associated to 2-year mortality

	95% CI	p
Age	1.032 (1.006-1.058)	0.015
Sex	1.335 (0.682-2.1615)	0.400
Charlson comorbidity index	1.266 (1.036-1.547)	0.021
Body mass index	0.934 (0.877-0.994)	0.032
Clinical frailty scale	1.406 (1.140-1.735)	0.001
Low handgrip strength	1.445 (0.697-2.995)	0.322

CI: Confidence interval

Discussion

This prospective cohort study demonstrated the independent effect of CFS and BMI on 2-year mortality prediction in hospitalized internal medicine patients. Whereas a higher CFS score was associated with a higher mortality risk, a lower BMI was associated with a higher mortality risk, regardless of age, sex, and CCI. In our study population, older patients had a higher CFS score. The rate of patients with low HGS and at risk of malnutrition was higher in the older group. This is not surprising as they are leading and challenging geriatric syndromes, especially for hospitalized older patients (12,26).

Recently, CFS has been widely used to predict adverse outcomes such as mortality in various settings (9). Although it was validated in geriatric patients (≥65 years), there are emerging studies suggesting its use at all ages (25). Welford et al. (27)

revealed that a higher CFS score was associated with a poor prognosis in hemato-oncology clinics. They supported the use of CFS in inpatients of any age. In another study, CFS was used in 18 years and older patients with cancer at an intensive care unit and was found to be associated with worse clinical outcomes among oncologic critically ill patients (28).

A multicenter retrospective cohort study with a median (IQR) age of 63.7 years (49.1-74.0 years) concluded that CFS predicted 1-year mortality well in critically ill patients (HR 1.26, 95% CI 1.21-1.31) after adjusting for confounders (29). A prospective multicenter cohort study was conducted in younger critically ill patients and supported the use of CFS in younger adults, not just in older adults (30). The results of our study are consistent with the literature. One-point increment in CFS was associated with 1.3-fold mortality risk in hospitalized internal medicine patients, regardless of age, sex, and comorbidities.

Another highlighted point of our study is the independent effect of BMI on mortality. We concluded that higher BMI scores were associated with lower mortality risk. A recently published, large sample size study conducted in geriatric medical departments presented the protective effect of BMI on mortality. They used standard BMI categories according to the World Health Organization in their study and emphasized the requirement of an ideal BMI for vulnerable groups (31). This result was similar to ours. In our study, the median (IQR) BMI scores were 25.8

Table 3. Multivariable Cox regression analysis associated to 2-year mortality

	Model-1		Model-2		Model-3		Model-4	
	CI 95%	p	CI 95%	p	CI 95%	p	CI 95%	p
Age	1.032 (1.006-1.058)	0.017	1.031 (1.003-1.060)	0.029	1.037 (1.011-1.063)	0.005	1.018 (0.987-1.050)	0.254
Sex	1.245 (0.635-2.441)	0.524	1.261 (0.643-2.474)	0.500	1.173 (0.597-2.305)	0.643	1.406 (0.672-2.945)	0.366
CCI			1.227 (1.002-1.504)	0.048	1.326 (1.076-1.634)	0.008	1.238 (0.991-1.547)	0.060
BMI					0.890 (0.830-0.954)	0.001	0.898 (0.840-0.961)	0.002
CFS							1.313 (1.002-1.719)	0.048

Model 1 age, sex; model 2 age, sex, CCI; model 3 age, sex, CCI, BMI; model 4 age, sex, CCI, BMI, CFS
 BMI: Body mass index, CCI: Charlson comorbidity index, CFS: Clinical frailty scale, CI: Confidence interval

(22.7-29.9) and 29.1 (25.5-32.1) for younger and older patients, respectively. On the other hand, we did not categorize patients according to BMI because the thresholds should be different for geriatric patients and patients living with frailty. In the light of these data, we evaluated BMI as a continuous variable and showed its effect on mortality irrespective of age, sex, CCI, and CFS. This striking point will provide a basis for future study designs. Kanenawa et al. (32) presented a study similar to ours. They determined the impact of CFS on 2-year mortality after hospitalization for heart failure, regardless of stratification based on age, sex, BMI, and left ventricular ejection fraction. Therefore, they suggested the use of CFS as a prognostic tool in clinical settings.

Study Limitations

There are some limitations to our study. First, CFS was not validated in younger patients. However, as there are so many studies supporting its use in younger patients, we used it for younger patients. Second, we evaluated BMI as a continuous variable and did not categorize it. We planned to investigate the effect of a 1-point change in BMI. Therefore, there are conflicting results regarding its use, especially for older adults. In this field, large sample size studies are needed, and the cut-off values for BMI should be assessed anew. In contrast, we highlighted the importance of using CFS and assessing BMI in hospitalized patients, regardless of age, sex, and CCI.

Conclusion

Higher CFS and lower BMI scores are independently associated with 2-year mortality in hospitalized internal medicine patients. Future comprehensive studies on the use of CFS in hospitalized patients and updating BMI cut-off values according to frailty and age categories are needed.

Ethics

Ethics Committee Approval: The study was approved by the Hacettepe University Non-Interventional Clinical Research Ethics Committee (decision number: 2019/07-28, date: 07.03.2019).

Informed Consent: Written informed consents were obtained from all participants.

Authorship Contributions

Surgical and Medical Practices: Y.Ö., M.G., M.K., M.H., Concept: Y.Ö., M.G., M.K., C.B., B.B.D., M.C., M.H., Design: Y.Ö., M.K., M.H., Data Collection or Processing: Y.Ö., M.G., S.C., M.K., B.B.D., M.C., M.H., Analysis or Interpretation: Y.Ö., A.O.B., M.G., S.C., M.E., C.B., B.B.D., M.C., M.H., Literature Search: Y.Ö., A.O.B., M.G., S.C., M.E., M.H., Writing: Y.Ö., A.O.B., M.G., S.C., M.K., M.E., C.B., M.H.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

- Clegg A, Young J, Iliffe S, Rikkert MO, Rockwood K. Frailty in elderly people. *Lancet* 2013;381:752-762.
- Hoogendijk EO, Afilalo J, Ensrud KE, Kowal P, Onder G, Fried LP. Frailty: implications for clinical practice and public health. *Lancet* 2019;394:1365-1375.
- Dent E, Martin FC, Bergman H, Woo J, Romero-Ortuno R, Walston JD. Management of frailty: opportunities, challenges, and future directions. *Lancet* 2019;394:1376-1386.
- Rezaei-Shahsavarloo Z, Atashzadeh-Shoorideh F, Gobbens RJJ, Ebadi A, Ghaedamini Harouni G. The impact of interventions on management of frailty in hospitalized frail older adults: a systematic review and meta-analysis. *BMC Geriatr* 2020;20:526.
- Morley JE, Vellas B, van Kan GA, Anker SD, Bauer JM, Bernabei R, Cesari M, Chumlea WC, Doehner W, Evans J, Fried LP, Guralnik JM, Katz PR, Malmstrom TK, McCarter RJ, Gutierrez Robledo LM, Rockwood K, von Haehling S, Vandewoude MF, Walston J. Frailty consensus: a call to action. *J Am Med Dir Assoc* 2013;14:392-397.
- Rolfson DB, Majumdar SR, Tsuyuki RT, Tahir A, Rockwood K. Validity and reliability of the Edmonton Frail Scale. *Age Ageing* 2006;35:526-529.
- Rockwood K, Song X, MacKnight C, Bergman H, Hogan DB, McDowell I, Mitnitski A. A global clinical measure of fitness and frailty in elderly people. *Cmaj* 2005;173:489-495.
- Rockwood K, Theou O. Using the Clinical Frailty Scale in Allocating Scarce Health Care Resources. *Can Geriatr J* 2020;23:210-215.

9. Church S, Rogers E, Rockwood K, Theou O. A scoping review of the Clinical Frailty Scale. *BMC Geriatr* 2020;20:393.
10. Cederholm T, Jensen GL, Correia MITD, Gonzalez MC, Fukushima R, Higashiguchi T, Baptista G, Barazzoni R, Blaauw R, Coats A, Crivelli A, Evans DC, Gramlich L, Fuchs-Tarlovsky V, Keller H, Llido L, Malone A, Mogensen KM, Morley JE, Muscaritoli M, Nyulasi I, Pirlich M, Pisprasert V, de van der Schueren MAE, Siltharm S, Singer P, Tappenden K, Velasco N, Waitzberg D, Yamwong P, Yu J, Van Gossum A, Compher C; GLIM Core Leadership Committee; GLIM Working Group. GLIM criteria for the diagnosis of malnutrition - A consensus report from the global clinical nutrition community. *Clin Nutr* 2019;38:1-9.
11. Cruz-Jentoft AJ, Kiesswetter E, Drey M, Sieber CC. Nutrition, frailty, and sarcopenia. *Aging Clin Exp Res* 2017;29:43-48.
12. Ligthart-Melis GC, Luiking YC, Kakourou A, Cederholm T, Maier AB, de van der Schueren MAE. Frailty, Sarcopenia, and Malnutrition Frequently (Co-) occur in Hospitalized Older Adults: A Systematic Review and Meta-analysis. *J Am Med Dir Assoc* 2020;21:1216-1228.
13. Riaz H, Khan MS, Siddiqi TJ, Usman MS, Shah N, Goyal A, Khan SS, Mookadam F, Krasuski RA, Ahmed H. Association Between Obesity and Cardiovascular Outcomes: A Systematic Review and Meta-analysis of Mendelian Randomization Studies. *JAMA Netw Open* 2018;1:e183788.
14. Li J, Li D, Wang X, Zhang L. The impact of body mass index on mortality rates of hip fracture patients: a systematic review and meta-analysis. *Osteoporos Int* 2022;33:1859-1869.
15. Kittiskulnam P, Johansen KL. The obesity paradox: A further consideration in dialysis patients. *Semin Dial* 2019;32:485-489.
16. Bosello O, Vanzo A. Obesity paradox and aging. *Eat Weight Disord* 2021;26:27-35.
17. Donini LM, Busetto L, Bischoff SC, Cederholm T, Ballesteros-Pomar MD, Batsis JA, Bauer JM, Boirie Y, Cruz-Jentoft AJ, Dicker D, Frara S, Frühbeck G, Genton L, Gepner Y, Giustina A, Gonzalez MC, Han HS, Heymsfield SB, Higashiguchi T, Laviano A, Lenzi A, Nyulasi I, Parrinello E, Poggiogalle E, Prado CM, Salvador J, Rolland Y, Santini F, Serlie MJ, Shi H, Sieber CC, Siervo M, Vettor R, Villareal DT, Volkert D, Yu J, Zamboni M, Barazzoni R. Definition and Diagnostic Criteria for Sarcopenic Obesity: ESPEN and EASO Consensus Statement. *Obes Facts* 2022;15:321-335.
18. Wang Z. Age-dependent decline of association between obesity and mortality: a systematic review and meta-analysis. *Obes Res Clin Pract* 2015;9:1-11.
19. Dramé M, Godaert L. The Obesity Paradox and Mortality in Older Adults: A Systematic Review. *Nutrients* 2023;15:1780.
20. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987;40:373-383.
21. Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyère O, Cederholm T, Cooper C, Landi F, Rolland Y, Sayer AA, Schneider SM, Sieber CC, Topinkova E, Vandewoude M, Visser M, Zamboni M; Writing Group for the European Working Group on Sarcopenia in Older People 2 (EWGSOP2), and the Extended Group for EWGSOP2. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing* 2019;48:16-31.
22. Ozturk Y, Deniz O, Coteli S, Unsal P, Dikmeer A, Burkuk S, Koca M, Cavusoglu C, Dogu BB, Cankurtaran M, Halil M. Global Leadership Initiative on Malnutrition criteria with different muscle assessments including muscle ultrasound with hospitalized internal medicine patients. *JPEN J Parenter Enteral Nutr* 2022;46:936-945.
23. Sarikaya D, Halil M, Kuyumcu ME, Kilic MK, Yesil Y, Kara O, Ozturk S, Gungor E, Karabulut E, Balam Yavuz B, Cankurtaran M, Ariogul S. Mini nutritional assessment test long and short form are valid screening tools in Turkish older adults. *Arch Gerontol Geriatr* 2015;61:56-60.
24. Bolayir B, Arık G, Yeşil Y, Kuyumcu ME, Varan HD, Kara Ö, Güngör AE, Yavuz BB, Cankurtaran M, Halil MG. Validation of Nutritional Risk Screening-2002 in a Hospitalized Adult Population. *Nutr Clin Pract* 2019;34:297-303.
25. Özsürekcı C, Balcı C, Kızırlanoğlu MC, Çalışkan H, Tuna Doğrul R, Ayçiçek GŞ, Sümer F, Karabulut E, Yavuz BB, Cankurtaran M, Halil MG. An important problem in an aging country: identifying the frailty via 9 Point Clinical Frailty Scale. *Acta Clin Belg* 2020;75:200-204.
26. Daly RM, Iuliano S, Fyfe JJ, Scott D, Kirk B, Thompson MQ, Dent E, Fetterplace K, Wright ORL, Lynch GS, Zanker J, Yu S, Kurrle S, Visvanathan R, Maier AB. Screening, Diagnosis and Management of Sarcopenia and Frailty in Hospitalized Older Adults: Recommendations from the Australian and New Zealand Society for Sarcopenia and Frailty Research (ANZSSFR) Expert Working Group. *J Nutr Health Aging* 2022;26:637-651.
27. Welford J, Rafferty R, Hunt K, Short D, Duncan L, Ward A, et al. The Clinical Frailty Scale can indicate prognosis and care requirements on discharge in oncology and haemato-oncology inpatients: A cohort study. *Eur J Cancer Care (Engl)* 2022;31:e13752.
28. Osatnik J, Matarrese A, Leone B, Cesar G, Kleinert M, Sosa F, Roberti J, Ivluch D. Frailty and clinical outcomes in critically ill patients with cancer: A cohort study. *J Geriatr Oncol* 2022;13:1156-1161.
29. Subramaniam A, Ueno R, Tiruvoipati R, Srikanth V, Bailey M, Pilcher D. Comparison of the predictive ability of clinical frailty scale and hospital frailty risk score to determine long-term survival in critically ill patients: a multicentre retrospective cohort study. *Crit Care* 2022;26:121.
30. Bagshaw M, Majumdar SR, Rolfson DB, Ibrahim Q, McDermid RC, Stelfox HT. A prospective multicenter cohort study of frailty in younger critically ill patients. *Crit Care* 2016;20:175.
31. Ryg J, Anru PL, Engberg H, Jorgensen MG, Masud T, Christensen K, Andersen-Ranberg K. Association of Body Mass Index With All-Cause Mortality in Acutely Hospitalized Older Patients. *J Am Med Dir Assoc* 2022;23:507-513.
32. Kanenawa K, Isotani A, Yamaji K, Nakamura M, Tanaka Y, Hirose-Inui K, Fujioka S, Mori S, Yano M, Ito S, Morinaga T, Fukunaga M, Hyodo M, Ando K. The impact of frailty according to Clinical Frailty Scale on clinical outcome in patients with heart failure. *ESC Heart Fail* 2021;8:1552-1561.